arXiv:1611.02902v2 [math.OC] 5 Feb 2017

Time-inconsistent stochastic control: solving the extended HJB system is a necessary condition for regular equilibria

Kristoffer Lindensjö*
Department of Mathematics, Stockholm University, SE-106 91 Stockholm, Sweden

February 7, 2017

Abstract

Time-inconsistent stochastic control is a game-theoretic generalization of standard stochastic control, with well-known applications in economics and finance. An important result of standard stochastic control is the characterization of the optimal value function as the solution to a PDE known as the HJB equation. Time-inconsistent stochastic control offers a similar possibility: Björk, Khapko and Murgoci (2016) [2] introduce a system of PDEs, the extended HJB system, and prove a verification theorem saying that *if* the extended HJB system has a solution then it is an equilibrium of a corresponding time-inconsistent stochastic control problem. In the present paper we show that a *regular* equilibrium is necessarily a solution to the extended HJB system. The model is that of a general (Markovian) Itô diffusion and general time-inconsistent running and terminal payoff functions.

Keywords: Dynamic inconsistency, Extended HJB system, Equilibrium, Hamilton-Jacobi-Bellman, Time inconsistent preferences, Time-inconsistent stochastic control

1 Introduction

Time-inconsistent stochastic control is a game theoretic generalization of standard stochastic control, based on the notion of Nash equilibrium. Time-inconsistent control problems have been studied in the finance and economics literature since at least the 1950s. The general mathematical theory of time-inconsistent stochastic control is, however, still highly incomplete, as Björk and Murgoci (2014) [5] write " *What has been lacking in the literature...is a...general theory of time-inconsistent stochastic control*".

We will now describe a simple standard (time-consistent) stochastic control problem in a somewhat unusual fashion and, as a contrast, introduce a time-inconsistent stochastic control problem.

*E-mail address: kristoffer.lindensjo@math.su.se.

A simple but standard stochastic control problem amounts to choosing a (deterministic) function $\mathbf{u} : [0,T] \times \mathbb{R}^n \to \mathbb{R}$ (known as a *Markovian feedback control law*) that maximizes

$$\tilde{J}(t,x,\mathbf{u}) = E_{t,x}[\tilde{F}(X_T^{\mathbf{u}})] \tag{1}$$

for each (t,x) in $[0,T] \times \mathbb{R}^n$, where $\tilde{F}$ is some nice real-valued function, often called *terminal payoff (function)*, and $X^{\mathbf{u}}$ is a controlled Itô diffusion

$$dX_s^{\mathbf{u}} = \mu(s, X_s^{\mathbf{u}}, \mathbf{u}(s, X_s^{\mathbf{u}}))ds + \sigma(s, X_s^{\mathbf{u}}, \mathbf{u}(s, X_s^{\mathbf{u}}))dW_s,$$

where W is a Wiener process and σ and μ are nice deterministic functions.

At first glance this optimization problem does not look well-posed since the optimal control (function) would apparently depend on the particular starting point (t,x). Fortunately, the problem turns out to be well-posed due to the *dynamic programming principle* (DPP), which can be heuristically formulated as follows: *if an optimal control for (1) exists for every fixed* (t,x) *then these controls do, in fact, coincide.* Thus, the problem of choosing a control $\mathbf{u}$ such that (1) is maximal *for each* (t,x) turns out to be well-posed.

The standard stochastic control theory cannot, however, satisfactorily handle the problem of choosing a control $\mathbf{u}$ such that it maximizes

$$J(t,x,\mathbf{u}) = E_{t,x}[F(X_T^{\mathbf{u}}, x)] \tag{2}$$

for each (t,x) in $[0,T] \times \mathbb{R}$, where F is a nice deterministic function which we call a *time-inconsistent terminal payoff (function)*. To see this, note that if t is current time and $X_t = x$, and if a control $\mathbf{u}$ is optimal at (t,x) then it will at any future point $(s, X_s^{\mathbf{u}})$ be optimal for the problem of maximizing $E_{s,X_s^{\mathbf{u}}}[F(X_T^{\mathbf{u}}, x)]$, according to the DPP. However, it will not generally be optimal for the problem of maximizing $E_{s,X_s^{\mathbf{u}}}[F(X_T^{\mathbf{u}}, X_s^{\mathbf{u}})]$ which is our task, cf. (2). This optimization problem is therefore equivalent to us having to deal with different terminal payoff functions at different points in time: at (t,x) we consider maximizing the conditional expected value (see (2)) based on the function $F(\cdot, x)$, but later, at for example $(s, X_s^{\mathbf{u}})$ we want to maximize the conditional expected value based on the currently unknown (random) function $F(\cdot, X_s^{\mathbf{u}})$. In other words: the set of preferences we have today are not the same as the set of preferences we have tomorrow. The optimization problem is therefore *time-inconsistent*, and it is in fact unclear what optimality for (2) even means. In other words, this problem is not well-posed.

In order to deal with the issue of time-inconsistency the literature has resorted to game theory and the notion of *Nash equilibrium*, not only for the setting above, but also more generally. Heuristically, the game-theoretic nature of the problem is due to the terminal payoff function changing over time, and we view the optimization problem as a game between agents (one agent for each time t) who optimize their own value function $E_{t,X_t^{\mathbf{u}}}[F(X_T^{\mathbf{u}}, X_t^{\mathbf{u}})]$, by choosing the control's (i.e. $\mathbf{u}$:s) value only at the time t. This idea is formalized by the introduction of an *equilibrium control* and a corresponding *equilibrium value function*, see Definition 3.2.

One of the main results of standard (time-consistent) stochastic control is the characterization of the optimal value function as the unique solution to a

certain non-linear partial differential equation known as the Hamilton-Jacobi-Bellman equation (HJB). This type of characterization was originally often performed under ad hoc assumptions regarding mainly sufficient differentiability (see Remark 8.2 for further comments). Another approach, which demands less regularity, is to consider *viscosity solutions*, see Remark 8.4. The characterization is often presented as two companion theorems (see e.g. [17, 19, 20, 24]) which in simplified versions can be phrased as follows: 1. *A necessary condition for two functions to be an optimal control and the optimal value function is that they solve the HJB*, and 2. *Solving the HJB is a sufficient condition for two functions to be an optimal control and the optimal value function* (this type of result is known as a *verification theorem*).

Naturally, we hope that time-inconsistent stochastic control problems offer a similar possibility. Indeed, an *extended HJB system* was proposed by Björk et al. around 2010, see [2].[1] The main result of [2] is a verification theorem which can be heuristically phrased as follows: *Solving the extended HJB system is a sufficient condition for being an equilibrium.*

The main contribution of the present paper can be summarized as follows.

- We study the general time-inconsistent stochastic control theory of Björk et al. [2] and show that solving the extended HJB system is a necessary condition for being a *regular* equilibrium, see Theorem 5.5 and the more general Theorem 6.3.
- We study a time-inconsistent version of the stochastic regulator problem as an example, see Section 7.

Remark 1.1 *We consider only regular equilibria. This means that we implicitly only consider models which satisfy certain conditions, mainly regarding the existence of an equilibrium control and differentiability (see Assumption 5.2 and Assumption 5.3). These assumptions, therefore, need to be verified for each particular model that is studied. We remark that the example studied in Section 7 has a regular equilibrium.*

1.1 Previous literature

Several particular time-inconsistent control problems were studied before the general theory studied in the present paper was first suggested by Björk et al. [2, 5]. The game-theoretic approach to time-inconsistency was first used by Strotz (1955) [23] when studying utility maximization problems. Other influential papers following a similar line in economics and finance are [15, 18, 22, 26]. These and other papers in mainly economics and finance have sparked a recent interest in time-inconsistent stochastic control from a more mathematical perspective. Early financial mathematics papers in time-inconsistent stochastic control include Ekeland and Lazrak (2006) [11] and Ekeland and Pirvu (2008) [12], who study the classic time-inconsistent finance problem of optimal consumption and investment under hyperbolic discounting. The ideas developed in these papers served as an inspiration to the general theory for time-inconsistent stochastic control proposed in [2, 5], which also include several interesting applications. A particular type of time-inconsistent stochastic control problem is

[1] Different versions of this paper have been available as preprints since at least 2010. The most cited version is [4].

mean-variance optimization (studied in e.g. optimal investment theory). Different versions of this problem have recently been studied by [6, 8, 21]. A numerical technique for mean-variance optimization problems is studied in [9].

1.1.1 Alternative approaches to time-inconsistent stochastic control

Yong (2012) [29] (see also [28]) and Wang and Wu (2015) [27] study time-inconsistent stochastic control, but they use a slightly different equilibrium construction than that of the present paper. They start with N agents each controlling the state process on a small time interval and then solve the equilibrium problem by using backward recursion, which is similar to how the problem is solved in discrete time in [5]. N is then sent to infinity. Moreover, in [27] the theory of FBSDEs is used to characterize the equilibrium (rather than with a PDE system). Djehiche and Huang (2015) [10] study a problem similar to that of the present paper, for which they characterize an equilibrium control (if it exists) via a stochastic maximum principle for general nonlinear diffusion models (i.e. as the solution to a particular SDE). Their main focus is, however, an equilibrium approach to mean field games.

2 The model

Let there be a stochastic basis $(\Omega, \mathcal{F}, P, \underline{\mathcal{F}})$ with a fixed time horizon $T < \infty$, where $\underline{\mathcal{F}}$ is the augmented filtration generated by a d-dimensional Wiener process W. We consider n-dimensional controlled diffusion processes

$$dX_s^{\mathbf{u}} = \mu(s, X_s^{\mathbf{u}}, \mathbf{u}(s, X_s^{\mathbf{u}}))ds + \sigma(s, X_s^{\mathbf{u}}, \mathbf{u}(s, X_s^{\mathbf{u}}))dW_s, \quad t \leq s \leq T, \quad X_t^{\mathbf{u}} = x \tag{3}$$

where,

- controls are measurable functions $\mathbf{u} : [0, T] \times \mathbb{R}^n \to U$ (known as Markovian feedback control laws), where $U \in \mathbb{R}^k$ is compact. We denote the set of such functions by $\mathbf{U}$.
- $\mu : [0, T] \times \mathbb{R}^n \times U \to \mathbb{R}^n$ and $\sigma : [0, T] \times \mathbb{R}^n \times U \to M(n, d)$ are bounded and continuous functions, where $M(n, d)$ denotes the set of real-valued matrices with dimension $n \times d$. Moreover, $\sigma(t, x, \mathbf{u}(t, x))$ is Lipschitz in $x \in \mathbb{R}^n$ and $\sigma(t, x, \mathbf{u}(t, x))\sigma(t, x, \mathbf{u}(t, x))^T$ is uniformly elliptic, for each $\mathbf{u} \in \mathbf{U}$, where ^{T} denotes transpose.

Definition 2.1 (Admissible control) *A control* $\mathbf{u}$ *is* admissible *if for any starting point* $(t, x) \in [0, T) \times \mathbb{R}^n$ *there exists a unique strong solution* $X^{\mathbf{u}}$ *to (3).* $X^{\mathbf{u}}$ *has continuous sample paths (by the definition of a strong solution).*

The time-inconsistent stochastic control problem, see Definition 3.1 below, is based on two different kinds of time-inconsistent terminal payoff functions F and G.

Definition 2.2 (Time-inconsistent terminal payoff functions)
$F : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}$ *is continuous and satisfies*

$$|F(x, y)| \leq C_0(1 + |x|^2) \tag{4}$$

for some constant C_0 (which may depend on y), where $|x|^2 := \sum_i x_i^2$.
$G : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}$ is twice continuously differentiable $G \in C^2(\mathbb{R}^n \times \mathbb{R}^n)$.

We will also use the following two functions.

Definition 2.3 (Auxiliary functions) *For $\mathbf{u} \in \mathbf{U}$ we define the function $f_{\mathbf{u}} : [0,T] \times \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}$ by*

$$f_{\mathbf{u}}(t,x,y) = E_{t,x}\left[F(X_T^{\mathbf{u}}, y)\right],$$

and the function $g_{\mathbf{u}} : [0,T] \times \times \mathbb{R}^n \to \mathbb{R}^n$ by

$$g_{\mathbf{u}}(t,x) = E_{t,x}\left[X_T^{\mathbf{u}}\right],$$

where $E_{t,x}[\cdot]$ denotes the expected value given the starting point (t,x) in (3).

Lemma 2.4

1. *Each control is an admissible control.*
2. *For any $\mathbf{u} \in \mathbf{U}$ and any $(t,x) \in [0,T] \times \mathbb{R}^n$, $E_{t,x}\left[(sup_{t \le s \le T}|X_s^{\mathbf{u}}|)^2\right] < C_1(1+|x|^2)$ for some constant C_1 which is independent of t and $\mathbf{u}$.*
3. *For any $\mathbf{u} \in \mathbf{U}$ and any $(t,x,y) \in [0,T] \times \mathbb{R}^n \times \mathbb{R}^n$, $|f_{\mathbf{u}}(t,x,y)| \le E_{t,x}[|F(X_T^{\mathbf{u}}, y)|] \le C_2(1+|x|^2)$ for some constant C_2 which is independent of t and $\mathbf{u}$ (but which may depend on y).*

Note that item 1. above implies that the set of controls $\mathbf{U}$ is actually also the set of admissible controls. Note also that item 2. implies that: for any $\mathbf{u} \in \mathbf{U}$ and any $(t,x) \in [0,T] \times \mathbb{R}^n$, $|g_{\mathbf{u}}(t,x)| < C_1(1+|x|^2)$ for some constant C_1 which is independent of t and $\mathbf{u}$.

Proof. For each function $\mathbf{u} \in \mathbf{U}$ and starting value (t,x) the SDE (3) has a unique strong solution $X^{\mathbf{u}}$, by a result found in e.g. [25] (and clarified in [16]). Each control is therefore admissible.

The integrability condition for $sup_{t \le s \le T}|X_s^{\mathbf{u}}|$ is a consequence of μ and σ being bounded (see e.g [17, ch. 5.3].). We remark that C_1 can be taken to depend only on n, T and the least upper bounds of $|\mu|$ and $|\sigma|$ (where $|\cdot|^2 = \Sigma_{i,j}(\cdot)^2$).

Using (4) and item 2. it is easy to see that

$$\begin{aligned} &|f_{\mathbf{u}}(t,x,y)| \le E_{t,x}\left[|F(X_T^{\mathbf{u}}, y)|\right] \le E_{t,x}\left[C_0(1+|X_T^{\mathbf{u}}|^2)\right] \\ &\le C_0 + C_0 E_{t,x}\left[|X_T^{\mathbf{u}}|^2\right] \le C_0 + C_0(C_1 + |x|^2) \le C_2(1+|x|^2) \end{aligned}$$

for some C_2 depending only on n, T, the least upper bounds of $|\mu|$ and $|\sigma|$ and C_0 (and therefore on y). ∎

We need some more notation.

- $\mathbf{A}^{\mathbf{u}}$ denotes the following indexed (by a control $\mathbf{u}$) differential operator

$$\mathbf{A}^{\mathbf{u}} = \frac{\partial}{\partial t} + \sum_{i=1}^{n} \mu_i(t,x,\mathbf{u}(t,x)) \frac{\partial}{\partial x_i} + \frac{1}{2} \sum_{i,j=1}^{n} \sigma\sigma_{ij}^T(t,x,\mathbf{u}(t,x)) \frac{\partial^2}{\partial x_i x_j}$$

  which operates on sufficiently differentiable functions $[0,T] \times \mathbb{R}^n \to \mathbb{R}$.

- $\mathbf{A^u}$ operates only on variables in parentheses. For any function $h(t,x,y) : [0,T] \times \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}$ we allow for the possibility of placing the third variable as a superscript $h^y(t,x) = h(t,x,y)$. This implies, for example, that $\mathbf{A^u}h^y(t,x)$ involves only derivatives with respect to the first and second variables t and x. Moreover, $\mathbf{A^u}h(t,x,x)$ is equal to $\mathbf{A^u}\bar{h}(t,x)$, where $\bar{h}(t,x) := h(t,x,x)$.

- $\mathbf{A^u}h^y(t,x)$ and $\mathbf{A^{u(t,x)}}h^y(t,x)$ denote the same object.

- If $g : [0,T] \times \mathbb{R}^n \to \mathbb{R}^n$, then, with obvious notation, $\mathbf{A^u}g(t,x) := (\mathbf{A^u}g_1(t,x), ..., \mathbf{A^u}g_n(t,x))^T$.

- We also need the indexed (by a control $\mathbf{u}$) differential operator $\mathbf{H^u}$ which, for sufficiently regular functions $G : \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}$ and $g : [0,T] \times \mathbb{R}^n \to \mathbb{R}^n$, is defined by
$$\mathbf{H}^{\mathbf{u}}_G g(t,x) = G_y(x, g(t,x)) \cdot \mathbf{A^u}g(t,x), \tag{5}$$
where $G_y(x,y)$ is the gradient of G with respect y for fixed x and $\cdot$ denotes the Euclidean inner product.

- We will use the following notation
$$G \diamond g(t,x) = G(x, g(t,x)). \tag{6}$$

Remark 2.5 *The expected values found throughout this paper are finite. To see this use arguments similar to those in the proof of Lemma 2.4.*

3 The time-inconsistent stochastic control problem and the definition of equilibrium

We consider the time-inconsistent terminal payoff functions F and G (Definition 2.2) and define the time-inconsistent value function as follows.

Definition 3.1 (Value function)

$$J(t,x,\mathbf{u}) := E_{t,x}\left[F(X^{\mathbf{u}}_T, x)\right] + G(x, E_{t,x}\left[X^{\mathbf{u}}_T\right]).$$

We remark that this definition is analogous to the one in [2]. Consider the problem of choosing a control $\mathbf{u}$ so that $J(t,x,\mathbf{u})$ is as large as possible for each (t,x). But now recall that this is not (yet) a well-posed problem, as we saw in the introduction. The first objective of our theory of time-inconsistent stochastic control is to make this a well-posed problem.

To this end, we introduce a game-theoretic notion of equilibrium very similar to that in [2], which was inspired by [11, 12]. The definition is a continuous time version of a *subgame perfect Nash equilibrium*.

Definition 3.2 (Equilibrium)

- $\hat{\mathbf{u}}$ *is an* **equilibrium control** *if it is an admissible control such that*
$$\liminf_{h \searrow 0} \frac{J(t,x,\hat{\mathbf{u}}) - J(t,x,\mathbf{u}_{t+h})}{h} \geq 0, \quad \forall (t,x) \in [0,T) \times \mathbb{R}^n, \tag{7}$$

for all $\mathbf{u}_{t+h} := \begin{cases} \mathbf{u}, & \text{on } [t,t+h]\times B[x] \\ \hat{\mathbf{u}}, & \text{on } \{[t,t+h]\times B[x]\}^c, \end{cases}$

where $\mathbf{u}\in\mathbf{U}$ *and* $B[x]$ *is some arbitrary* n*-dimensional ball centered at* x*.*

- *If* $\hat{\mathbf{u}}$ *is an equilibrium control then the corresponding* **equilibrium value function** *is defined by* $V_{\hat{\mathbf{u}}}(t,x) = J(t,x,\hat{\mathbf{u}})$*.*
- *An* **equilibrium** *is a quadruple consisting of an equilibrium control, the corresponding equilibrium value function and the corresponding auxiliary functions* $(\hat{\mathbf{u}}, V_{\hat{\mathbf{u}}}(t,x), f_{\hat{\mathbf{u}}}(t,x,y), g_{\hat{\mathbf{u}}}(t,x))$*.*

The equilibrium value function $V_{\hat{\mathbf{u}}}(t,x)$ is indexed by the equilibrium control since several equilibrium controls may exist and correspond to different equilibrium value functions. Note that $\mathbf{u}_{t+h}$ clearly depends on x and $|B[x]|$, and also on t and h rather than $t+h$. The fact that our notation does not indicate this should not cause any confusion.

Definition 2.3, Definition 3.1, Definition 3.2 and (6) imply the following identities

$$V_{\hat{\mathbf{u}}}(t,x) = J(t,x,\hat{\mathbf{u}}) = f_{\hat{\mathbf{u}}}(t,x,x) + G\diamond g_{\hat{\mathbf{u}}}(t,x). \tag{8}$$

Remark 3.3 (Interpretation of $\mathbf{u}_{t+h}$) *If we at* (t,x) *consider* $\mathbf{u}_{t+h}$ *then we consider an alternative control* $\mathbf{u}$ *(compared to* $\hat{\mathbf{u}}$*) which is to be used until either 1. the short time period* $h>0$ *has passed, or 2. the controlled process* $X^{\mathbf{u}}$ *has left the ball* $B[x]$*.*

Remark 3.4 (Interpretation of equilibrium) *We can think of the agents in this game as different time versions of a person who faces the control problem corresponding to the value function* $J(t,x,\mathbf{u})$*. The interpretation of an equilibrium control is that it is a control which every agent wants to use if every agent after her (i.e. every future version of the person) uses that control. Heuristically, we can for small* h *think of* $J(t,x,\hat{\mathbf{u}}) - J(t,x,\mathbf{u}_{t+h})$ *as the expected payoff an agent at time* t *(the current "self") obtains when choosing the equilibrium control* $\hat{\mathbf{u}}$ *minus the expected payoff she would obtain if she chose the alternative control* $\mathbf{u}$*, provided that* $\hat{\mathbf{u}}$ *is used when she is no longer able to choose the control. Note that* $J(t,x,\hat{\mathbf{u}}) - J(t,x,\mathbf{u}_{t+h})$ *would approach zero when sending* h *to zero, and we therefore normalize this difference in (7).*

Remark 3.5 $\mathbf{u}_{t+h}$ *is bounded and measurable (cf. pasting of measurable functions) and is therefore in* $\mathbf{U}$*. This remark also applies to all controls we introduce later.*

4 The extended HJB system

Here we define the extended HJB system corresponding to the time-inconsistent payoff functions F and G i Definition 2.2. The definition is equivalent to the one in [2].

Definition 4.1 (The extended HJB system) *For* $(t,x,y) \in [0,T) \times \mathbb{R}^n \times \mathbb{R}^n$,

$$\begin{aligned}
\mathcal{A}^{\bar{\mathbf{u}}} f^y(t,x) = 0, \quad f^y(T,x) = F(x,y) \\
\mathcal{A}^{\bar{\mathbf{u}}} g(t,x) = 0, \quad g(T,x) = x \\
\sup_{u \in U} \left\{ \mathbf{A}^u V(t,x) - \mathbf{A}^u f(t,x,x) + \mathbf{A}^u f^x(t,x) - \mathbf{A}^u G \diamond g(t,x) + \mathbf{H}^u_G g(t,x) \right\} = 0 \\
V(T,x) = F(x,x) + G(x,x)
\end{aligned}$$

where $\bar{\mathbf{u}}$ *satisfies*

$$\bar{\mathbf{u}}(t,x) \in \arg\sup_{u \in U} \left\{ \mathbf{A}^u V(t,x) - \mathbf{A}^u f(t,x,x) + \mathbf{A}^u f^x(t,x) - \mathbf{A}^u G \diamond g(t,x) + \mathbf{H}^u_G g(t,x) \right\}. \tag{9}$$

A solution to the extended HJB system is a quadruple $(\bar{\mathbf{u}}, V(t,x), f(t,x,y), g(t,x))$ where $\bar{\mathbf{u}} : [0,T] \times \mathbb{R}^n \to U, V : [0,T] \times \mathbb{R}^n \to \mathbb{R}, f : [0,T] \times \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}$ and $g : [0,T] \times \mathbb{R}^n \to \mathbb{R}^n$. In the next section we will see that a regular equilibrium (Definition 5.1) is classical solution to the extended HJB system. By a classical solution we mean that the functions in the equilibrium are sufficiently differentiable for the derivatives of the extended HJB system to exist in the usual sense.

Remark 4.2 *The maximization in (9) should be construed as follows. For a fixed point* (t,x) *and any, sufficiently differentiable, fixed functions* $V(t,x)$, $f(t,x,y)$ *and* $g(t,x)$ *the maximization procedure regards*

$$\mathbf{A}^u V(t,x) - \mathbf{A}^u f(t,x,x) + \mathbf{A}^u f^x(t,x) - \mathbf{A}^u G \diamond g(t,x) + \mathbf{H}^u_G g(t,x) \tag{10}$$

with respect to $u \in U$*. Note that the definitions in (5) and (6) reveal exactly how the last two objects in (10) depend on* $g(t,x)$*. Also note that* $\mathbf{A}^u f^x(t,x)$ *involves derivatives only with respect to* t *and the first* x *in* $f(t,x,x)$*.*

Remark 4.3 *The two first rows of the extended HJB system correspond to two Kolmogorov backward equations indexed by* $\bar{\mathbf{u}}$*, we call these rows the* **Kolmogorov part***. More precisely, the first row corresponds to a family of Kolmogorov backward equations, one for each* $y \in \mathbb{R}^n$*, while the second row is a standard Kolmogorov backward equation (whose solution is a function* $[0,T] \times \mathbb{R}^n \to \mathbb{R}^n$*). The third and fourth row, we call these rows the* **HJB part***, consist of a standard HJB equation, but with the non-standard feature that it includes derivatives of the solutions to the Kolmogorov part.*

It may shed some light on the extended HJB system to consider the following schedule. Start by guessing a solution, i.e. any four functions $\bar{\mathbf{u}}, V(t,x), f(t,x,y)$ and $g(t,x)$ which have the right dimensions and are sufficiently differentiable, and then

1. check that the boundary conditions $V(T,x) = F(x,x)+G(x,x), f(T,x,y) = F(x,y)$ and $g(T,x) = x$ are satisfied,
2. check that $f(t,x,y)$ and $g(t,x)$ satisfy the Kolmogorov part with the differential operator indexed with $\bar{\mathbf{u}}$,
3. for each (t,x), check that at least one maximizer of (10) exists, produces the maximal value 0, and is equal to the value $\bar{\mathbf{u}}(t,x)$.

If the guessed functions $\bar{\mathbf{u}}, V(t,x), f(t,x,y)$ and $g(t,x)$ satisfy 1,2 and 3, then they are a solution to the extended HJB system.

5 A regular equilibrium necessarily solves the extended HJB system

Let us first define what we mean by a regular equilibrium.

Definition 5.1 (Regular equilibrium) *An equilibrium (Definition 3.2) $(\hat{\mathbf{u}}, V(t,x), f_{\hat{\mathbf{u}}}(t,x,y), g_{\hat{\mathbf{u}}}(t,x))$ is said to be regular if Assumption 5.2 and Assumption 5.3 below are satisfied.*

Assumption 5.2 (Right-continuous equilibrium control) *There exists a right-continuous function $\hat{\mathbf{u}} : [0,T] \times \mathbb{R}^n \to U$ such that (7) is satisfied.*

Assumption 5.3 (Differentiability) *The auxiliary functions and the equilibrium value function corresponding to $\hat{\mathbf{u}}$ satisfy the following conditions.*

1. *$f^y_{\hat{\mathbf{u}}}(t,x) \in C^{1,2}([0,T) \times \mathbb{R}^n) \cap C([0,T] \times \mathbb{R}^n)$ for each fixed $y \in \mathbb{R}^n$.*
2. *The elements of $(g_{\hat{\mathbf{u}},1}(t,x), ..., g_{\hat{\mathbf{u}},n}(t,x))^T := g_{\hat{\mathbf{u}}}(t,x)$ satisfy $g_{\hat{\mathbf{u}},i}(t,x) \in C^{1,2}([0,T) \times \mathbb{R}^n) \cap C([0,T] \times \mathbb{R}^n)$, $i = 1, ..., n$.*
3. *The derivatives $\frac{\partial f_{\hat{\mathbf{u}}}(t,x,x)}{\partial t}$, $\frac{\partial f_{\hat{\mathbf{u}}}(t,x,x)}{\partial x_i}$, $\frac{\partial^2 f_{\hat{\mathbf{u}}}(t,x,x)}{\partial x_i \partial x_j}$ exist for each $(t,x) \in [0,T) \times \mathbb{R}^n$ and $i,j = 1, ..., n$.*

Remark 5.4 *We consider only regular equilibria. This means that we implicitly only consider models (i.e. σ, μ, U, F and G) which are such that Assumption 5.2 and Assumption 5.3 are satisfied. Thus, these assumptions need to be verified for each particular model that is studied. See Section 7 for a concrete example with a regular equilibrium.*

We are now ready to present our main result.

Theorem 5.5 *A regular equilibrium $(\hat{\mathbf{u}}, V_{\hat{\mathbf{u}}}(t,x), f_{\hat{\mathbf{u}}}(t,x,y), g_{\hat{\mathbf{u}}}(t,x))$ necessarily solves the extended HJB system (Definition 4.1). In particular, if a regular equilibrium exists then the extended HJB system has a solution.*

Proof. The last assertion of the theorem follows directly from the first. The proof comprises 7 parts.

Part 1. From Definition 2.3, Definition 3.1 and Definition 3.2 it directly follows that $V_{\hat{\mathbf{u}}}(T,x) = F(x,x) + G(x,x)$, $f_{\hat{\mathbf{u}}}(T,x,y) = F(x,y)$ and $g_{\hat{\mathbf{u}}}(T,x) = x$. The equilibrium therefore satisfies the boundary conditions of the extended HJB system.

Part 2. Here we will show that the auxiliary function

$$f^y_{\hat{\mathbf{u}}}(t,x) = E_{t,x}[F(X^{\hat{\mathbf{u}}}_T, y)] \tag{11}$$

solves the first part of the Kolmogorov part of the extended HJB system. We remark that this corresponds to a version of the Feynman-Kac formula.

Let (t,x,y) be an arbitrary point in $[0,T) \times \mathbb{R}^n \times \mathbb{R}^n$. Consider the sequence of stopping times given by

$$\tau_n := \inf\left\{s > t : (s, X^{\hat{\mathbf{u}}}_s) \notin [t, t+h_n) \times B[x]\right\} \wedge T \tag{12}$$

where $B[x]$ is an arbitrary ball centered at x and $\{h_n\}$ is an arbitrary positive and decreasing sequence satisfying $\lim_{n\to\infty} h_n = 0$. The hitting time (12) is a stopping time, see e.g. [1, 17]. It follows directly from (12) that $X_s^{\hat{\mathbf{u}}}$ is bounded on the stochastic interval $[t, \tau_n]$. Note also that for almost all outcomes $\omega \in \Omega$ there exists an $N(\omega)$ such that $\tau_n = t + h_n$ for $n \geq N(\omega)$, so that $\tau_n \to t$ in the same way as $t + h_n \to t$ as $n \to \infty$ a.s.

Now, view $f_{\hat{\mathbf{u}}}^y(s, X_s^{\hat{\mathbf{u}}})_{s\in[t,T]}$ as a stochastic process and use Itô's formula (the necessary differentiability is provided by Assumption 5.3) to see that

$$\begin{aligned} f_{\hat{\mathbf{u}}}^y(t,x) \quad &= E_{t,x}\left[f_{\hat{\mathbf{u}}}^y(\tau_n, X_{\tau_n}^{\hat{\mathbf{u}}}) - \int_t^{\tau_n} \mathbf{A}^{\hat{\mathbf{u}}} f_{\hat{\mathbf{u}}}^y(s, X_s^{\hat{\mathbf{u}}})ds\right] \\ &\quad - E_{t,x}\left[\int_t^{\tau_n} \sum_{i=1}^n \sigma_i(s, X_s^{\hat{\mathbf{u}}}, \hat{\mathbf{u}}(s, X_s^{\hat{\mathbf{u}}})) \frac{\partial f_{\hat{\mathbf{u}}}^y(s, X_s^{\hat{\mathbf{u}}})}{\partial x_i} dW_s\right] \\ &= E_{t,x}\left[f_{\hat{\mathbf{u}}}^y(\tau_n, X_{\tau_n}^{\hat{\mathbf{u}}}) - \int_t^{\tau_n} \mathbf{A}^{\hat{\mathbf{u}}} f_{\hat{\mathbf{u}}}^y(s, X_s^{\hat{\mathbf{u}}})ds\right] \end{aligned} \tag{13}$$

where σ_i is the i:th row of σ and where the expected value of the Itô integral is zero since its integrand is bounded (which follows from the facts that $X_s^{\hat{\mathbf{u}}}$ is, for each n, bounded on the stochastic interval $[t, \tau_n]$, σ is bounded, and the derivatives $\frac{\partial f_{\hat{\mathbf{u}}}^y(s, X_s^{\hat{\mathbf{u}}})}{\partial x_i}$ satisfy Assumption 5.3).

Now, use (11) and then the tower property (Lemma 2.4 ensures that $F(y, X_T^{\hat{\mathbf{u}}})$ is integrable) to see that

$$E_{t,x}\left[f_{\hat{\mathbf{u}}}^y(\tau_n, X_{\tau_n}^{\hat{\mathbf{u}}})\right] = E_{t,x}\left[E_{\tau_n, X_{\tau_n}^{\hat{\mathbf{u}}}}\left[F(X_T^{\hat{\mathbf{u}}}, y)\right]\right] = E_{t,x}\left[F(X_T^{\hat{\mathbf{u}}}, y)\right] = f_{\hat{\mathbf{u}}}^y(t,x),$$

which with (13) implies, for each n, that

$$E_{t,x}\left[\frac{\int_t^{\tau_n} \mathbf{A}^{\hat{\mathbf{u}}} f_{\hat{\mathbf{u}}}^y(s, X_s^{\hat{\mathbf{u}}})ds}{h_n}\right] = 0. \tag{14}$$

Consider now the sequence of random variables given by

$$\frac{\int_t^{\tau_n} \mathbf{A}^{\hat{\mathbf{u}}} f_{\hat{\mathbf{u}}}^y(s, X_s^{\hat{\mathbf{u}}})ds}{h_n}. \tag{15}$$

Note that the integrand in (15) is bounded and has right-continuous trajectories on $[t, \tau_n]$, for each n.[2] It follows that we may, for each fixed outcome $\omega \in \Omega$, view the numerator of (15) as a deterministic Lebesgue integral of a one-dimensional right-continuous bounded deterministic function (in s).

Recall that $\tau_n \to t$ in the same way as $t + h_n \to t$ as $n \to \infty$ a.s. It follows from Lebesgue's differentiation theorem that sending $n \to \infty$ in (15) gives us

$$\lim_{n\to\infty} \frac{\int_t^{\tau_n} \mathbf{A}^{\hat{\mathbf{u}}} f_{\hat{\mathbf{u}}}^y(s, X_s^{\hat{\mathbf{u}}})ds}{h_n} = \mathbf{A}^{\hat{\mathbf{u}}} f_{\hat{\mathbf{u}}}^y(t,x) \quad a.s. \tag{16}$$

[2] To see this use that the integrand of (15) is

$$\mathbf{A}^{\hat{\mathbf{u}}} f_{\hat{\mathbf{u}}}^y(s, X_s^{\hat{\mathbf{u}}}) = \frac{\partial f_{\hat{\mathbf{u}}}^y(s, X_s^{\hat{\mathbf{u}}})}{\partial s} + \sum_{i=1}^n \mu_i(s, X_s^{\hat{\mathbf{u}}}, \hat{\mathbf{u}}(s, X_s^{\hat{\mathbf{u}}})) \frac{\partial f_{\hat{\mathbf{u}}}^y(s, X_s^{\hat{\mathbf{u}}})}{\partial x_i} + \frac{1}{2} \sum_{i,j=1}^n \sigma\sigma_{ij}^T(s, X_s^{\hat{\mathbf{u}}}, \hat{\mathbf{u}}(s, X_s^{\hat{\mathbf{u}}})) \frac{\partial^2 f_{\hat{\mathbf{u}}}^y(s, X_s^{\hat{\mathbf{u}}})}{\partial x_i x_j},$$

where $\mu(s, x, \hat{\mathbf{u}}(s,x))$ and $\sigma(s, x, \hat{\mathbf{u}}(s,x))$ are bounded and right-continuous (Section 2 and Assumption 5.2), and the derivatives satisfy Assumption 5.3. Moreover, recall that $X^{\hat{\mathbf{u}}}$ has continuous trajectories and that $X_s^{\hat{\mathbf{u}}}$ is bounded on $[t, \tau_n]$.

The random variables in (15) are bounded by a constant (uniformly in n[3]) and they correspond to an a.s. convergent sequence (cf. (16)) and we may therefore use dominated convergence when sending $n \to \infty$ in (14), which with (16) gives us

$$0 = \lim_{n\to\infty} E_{t,x}\left[\frac{\int_t^{\tau_n} \mathbf{A}^{\hat{\mathbf{u}}} f^y_{\hat{\mathbf{u}}}(s, X^{\hat{\mathbf{u}}}_s)ds}{h_n}\right] = E_{t,x}\left[\lim_{n\to\infty} \frac{\int_t^{\tau_n} \mathbf{A}^{\hat{\mathbf{u}}} f^y_{\hat{\mathbf{u}}}(s, X^{\hat{\mathbf{u}}}_s)ds}{h_n}\right]$$
$$= E_{t,x}\left[\mathbf{A}^{\hat{\mathbf{u}}} f^y_{\hat{\mathbf{u}}}(t,x)\right] = \mathbf{A}^{\hat{\mathbf{u}}} f^y_{\hat{\mathbf{u}}}(t,x)$$

which, since (t,x,y) was arbitrary, means that $f_{\hat{\mathbf{u}}}(t,x,y)$ solves the first part of the Kolmogorov part of the extended HJB system.

Part 3. Using arguments analogous to those in Part 2 it can be shown that the auxiliary function

$$g_{\hat{\mathbf{u}}}(t,x) = E_{t,x}[X^{\hat{\mathbf{u}}}_T]$$

solves the second part of the Kolmogorov part of the extended HJB system.

Part 4. The conclusion of Part 3 means that $0 = \mathbf{A}^{\hat{\mathbf{u}}} g_{\hat{\mathbf{u}}}(t,x)$, which implies that

$$\mathbf{H}^{\hat{\mathbf{u}}}_G g_{\hat{\mathbf{u}}}(t,x) = G_y(x, g_{\hat{\mathbf{u}}}(t,x)) \cdot \mathbf{A}^{\hat{\mathbf{u}}} g_{\hat{\mathbf{u}}}(t,x) = 0 \tag{17}$$

for all $(t,x) \in [0,T) \times \mathbb{R}^n$. Using (8) we obtain $\mathbf{A}^{\hat{\mathbf{u}}} V_{\hat{\mathbf{u}}}(t,x) = \mathbf{A}^{\hat{\mathbf{u}}} f_{\hat{\mathbf{u}}}(t,x,x) + \mathbf{A}^{\hat{\mathbf{u}}} G \diamond g_{\hat{\mathbf{u}}}(t,x) = 0$ (the necessary differentiability is provided by Definition 2.2 and Assumption 5.3), which with (17) and the conclusion of Part 2 gives us

$$\mathbf{A}^{\hat{\mathbf{u}}} V_{\hat{\mathbf{u}}}(t,x) - \mathbf{A}^{\hat{\mathbf{u}}} f_{\hat{\mathbf{u}}}(t,x,x) + \mathbf{A}^{\hat{\mathbf{u}}} f^x_{\hat{\mathbf{u}}}(t,x) - \mathbf{A}^{\hat{\mathbf{u}}} G \diamond g_{\hat{\mathbf{u}}}(t,x) + \mathbf{H}^{\hat{\mathbf{u}}}_G g_{\hat{\mathbf{u}}}(t,x) = 0 \tag{18}$$

for all $(t,x) \in [0,T) \times \mathbb{R}^n$.

The conclusions of Part 1, Part 2, Part 3 and Part 4 imply that the regular equilibrium $(\hat{\mathbf{u}}, V_{\hat{\mathbf{u}}}(t,x), f_{\hat{\mathbf{u}}}(t,x,y), g_{\hat{\mathbf{u}}}(t,x))$ can be seen as a *candidate solution* to the extended HJB system in the sense that the only thing we now have left to do is to show that our equilibrium control $\hat{\mathbf{u}}$ is maximal in (18), for each (t,x). This is what we will do in the rest of the proof.

Part 5: Consider an arbitrary point $(t,x,u) \in [0,T) \times \mathbb{R}^n \times U$ and the sequence of stopping times given by

$$\tilde{\tau}_n := \inf\{s > t : (s, X^u_s) \notin [t, t+h_n) \times B[x]\} \wedge T. \tag{19}$$

Let $u_{\tilde{\tau}_n}$ denote a control that is equal to the arbitrary constant u on $[t, \tilde{\tau}_n]$ and equal to the equilibrium control $\hat{\mathbf{u}}$ outside of this interval. Using (19), we note that $u_{\tilde{\tau}_n}$ is equal to u on $[t, t+h_n] \times B[x]$ and equal to $\hat{\mathbf{u}}$ on $\{[t, t+h_n] \times B[x]\}^c$, and hence that $u_{\tilde{\tau}_n}$ is, for each n, a (deterministic) function of the type $\mathbf{u}_{t+h}$ in Definition 3.2, specifically $u_{\tilde{\tau}_n} = u_{t+h_n}$.

Note that

$$E_{\tilde{\tau}_n, X^{u_{\tilde{\tau}_n}}_{\tilde{\tau}_n}}[F(X^{\hat{\mathbf{u}}}_T, x)] = E_{\tilde{\tau}_n, X^{u_{\tilde{\tau}_n}}_{\tilde{\tau}_n}}[F(X^{u_{\tilde{\tau}_n}}_T, x)] \quad a.s. \tag{20}$$

[3]The boundedness for a specific n is obvious (since $X^{\hat{\mathbf{u}}}_s$ is bounded by a constant on $[t, \tau_n]$, cf. (12), the derivatives of $f^y_{\hat{\mathbf{u}}}(t,x)$ satisfy Assumption 5.3, and μ and σ are bounded (Section 2)). The uniform boundedness can explained as follows: for (almost all) fixed ω (and our fixed (t,x,y) and $\hat{\mathbf{u}}$), the integrals in (15) correspond to a convergent sequence (in $\mathbb{R}$, with index n) where each element is bounded by a constant $M_n \in \mathbb{R}$ which does not depend on ω (cf. (12)), where the sequence $\{M_n\}$ can be taken to converge (cf. the right side of (16)). It is easy to see that there for such a sequence exists a constant M dominating each element of the sequence (where M then also does not depend on ω), which is what we claimed.

since the starting values of the two expected values are both $(\tilde{\tau}_n, X_{\tilde{\tau}_n}^{u_{\tilde{\tau}_n}})$, and since the controls (i.e. $u_{\tilde{\tau}_n}$ and $\hat{\mathbf{u}}$) are equal after the starting time, i.e. on $[\tilde{\tau}_n, T]$ except at exactly the starting time $\tilde{\tau}_n$. The only difference between the controls is thus on a set with Lebesgue measure zero, and the expected values in (20) therefore coincide.

Now use Itô's formula, in the same way as in Part 2, to see that

$$E_{t,x}\left[f_{\hat{\mathbf{u}}}^x(\tilde{\tau}_n, X_{\tilde{\tau}_n}^{u_{\tilde{\tau}_n}})\right] = f_{\hat{\mathbf{u}}}^x(t,x) + E_{t,x}\left[\int_t^{\tilde{\tau}_n} \mathbf{A}^{u_{\tilde{\tau}_n}} f_{\hat{\mathbf{u}}}^x(s, X_s^{u_{\tilde{\tau}_n}})ds\right]. \tag{21}$$

The integral limits in (21) are t and $\tilde{\tau}_n$ and the control $u_{\tilde{\tau}_n}$ is equal to u on $[t, \tilde{\tau}_n]$. It follows that $\mathbf{A}^{u_{\tilde{\tau}_n}} = \mathbf{A}^u$ and $X_s^{u_{\tilde{\tau}_n}} = X_s^u$ a.s. on $[t, \tilde{\tau}_n]$. We may therefore replace the superscript in the differential operator and in the controlled process in the integral in (21) with u. Using this replacement and (20) we rewrite (21) as

$$\begin{aligned}
&f_{\hat{\mathbf{u}}}^x(t,x) + E_{t,x}\left[\int_t^{\tilde{\tau}_n} \mathbf{A}^u f_{\hat{\mathbf{u}}}^x(s, X_s^u)ds\right] = E_{t,x}\left[f_{\hat{\mathbf{u}}}^x(\tilde{\tau}_n, X_{\tilde{\tau}_n}^{u_{\tilde{\tau}_n}})\right] \\
&= E_{t,x}\left[E_{\tilde{\tau}_n, X_{\tilde{\tau}_n}^{u_{\tilde{\tau}_n}}}\left[F(X_T^{\hat{\mathbf{u}}}, x)\right]\right] = E_{t,x}\left[E_{\tilde{\tau}_n, X_{\tilde{\tau}_n}^{u_{\tilde{\tau}_n}}}\left[F(X_T^{u_{\tilde{\tau}_n}}, x)\right]\right] \\
&= E_{t,x}\left[F(X_T^{u_{\tilde{\tau}_n}}, x)\right] = f_{u_{\tilde{\tau}_n}}(t,x,x).
\end{aligned}$$

Using dominated convergence and Lebesgue's differentiation theorem (which we may use for reasons analogous to those in Part 2) we thus obtain

$$\begin{aligned}
&\lim_{n\to\infty} \frac{f_{\hat{\mathbf{u}}}(t,x,x) - f_{u_{\tilde{\tau}_n}}(t,x,x)}{h_n} \\
&= \lim_{n\to\infty} \frac{-E_{t,x}\left[\int_t^{\tilde{\tau}_n} \mathbf{A}^u f_{\hat{\mathbf{u}}}^x(s, X_s^u)ds\right]}{h_n} \\
&= E_{t,x}\left[\lim_{n\to\infty} \frac{-\int_t^{\tilde{\tau}_n} \mathbf{A}^u f_{\hat{\mathbf{u}}}^x(s, X_s^u)ds}{h_n}\right] \\
&= -\mathbf{A}^u f_{\hat{\mathbf{u}}}^x(t,x).
\end{aligned} \tag{22}$$

Part 6: Consider an arbitrary point $(t,x,u) \in [0,T) \times \mathbb{R}^n \times U$ and the stopping times in (19). Using arguments analogous to those in Part 6 and standard differentiation rules we obtain

$$\begin{aligned}
&\lim_{n\to\infty} \frac{G \diamond g_{\hat{\mathbf{u}}}(t,x) - G \diamond g_{u_{\tilde{\tau}_n}}(t,x)}{h_n} \\
&= -\lim_{n\to\infty} \frac{G(x, g_{u_{\tilde{\tau}_n}}(t,x)) - G(x, g_{\hat{\mathbf{u}}}(t,x))}{h_n} \\
&= -\lim_{n\to\infty} \frac{G(x, E_{t,x}[X_T^{u_{\tilde{\tau}_n}}]) - G(x, g_{\hat{\mathbf{u}}}(t,x))}{h_n} \\
&= -\lim_{n\to\infty} \frac{G(x, E_{t,x}\left[E_{\tilde{\tau}_n, X_{\tilde{\tau}_n}^{u_{\tilde{\tau}_n}}}[X_T^{u_{\tilde{\tau}_n}}]\right]) - G(x, g_{\hat{\mathbf{u}}}(t,x))}{h_n} \\
&= -\lim_{n\to\infty} \frac{G\left(x, E_{t,x}\left[E_{\tilde{\tau}_n, X_{\tilde{\tau}_n}^{u_{\tilde{\tau}_n}}}[X_T^{\hat{\mathbf{u}}}]\right]\right) - G(x, g_{\hat{\mathbf{u}}}(t,x))}{h_n}
\end{aligned}$$

$$
\begin{aligned}
&= -\lim_{n\to\infty} \frac{G\left(x, E_{t,x}\left[g_{\hat{\mathbf{u}}}(\tilde{\tau}_n, X^{u_{\tilde{\tau}_n}}_{\tilde{\tau}_n})\right]\right) - G(x, g_{\hat{\mathbf{u}}}(t,x))}{h_n} \\
&= -\lim_{n\to\infty} \frac{G\left(x, g_{\hat{\mathbf{u}}}(t,x) + E_{t,x}\left[\int_t^{\tilde{\tau}_n} \mathbf{A}^u g_{\hat{\mathbf{u}}}(s, X^u_s) ds\right]\right) - G(x, g_{\hat{\mathbf{u}}}(t,x))}{h_n} \\
&= -G_y(x, g_{\hat{\mathbf{u}}}(t,x)) \cdot \mathbf{A}^u g_{\hat{\mathbf{u}}}(s,x) \\
&= -\mathbf{H}^u_G g_{\hat{\mathbf{u}}}(t,x) \qquad (23)
\end{aligned}
$$

where the last equality is true by definition, see (5).

Part 7: Let us verify the inequality and equalities below. The inequality follows from the assumption that $\hat{\mathbf{u}}$ is an equilibrium control (Definition 3.2). The first equality follows the fact that $u_{\tilde{\tau}_n} = u_{t+h_n}$ (see the arguments directly after (19)), which implies that we may replace u_{t+h} with $u_{\tilde{\tau}_n}$, h with h_n and $h \searrow 0$ with $n \to \infty$. The second equality follows from the definition of $J(t,x,\mathbf{u})$. The third equality follow from (22) and (23). The last equality follows from (8).

$$
\begin{aligned}
0 \quad &\geq -\lim_{h\searrow 0} \frac{J(t,x,\hat{\mathbf{u}}) - J(t,x,u_{t+h})}{h} \\
&= -\lim_{n\to\infty} \frac{J(t,x,\hat{\mathbf{u}}) - J(t,x,u_{\tilde{\tau}_n})}{h_n} \\
&= -\lim_{n\to\infty} \frac{f_{\hat{\mathbf{u}}}(t,x,x) + G \diamond g_{\hat{\mathbf{u}}}(t,x) - (f_{u_{\tilde{\tau}_n}}(t,x,x) + G \diamond g_{u_{\tilde{\tau}_n}})}{h_n} \\
&= -(-\mathbf{A}^u f^x_{\hat{\mathbf{u}}}(t,x) - \mathbf{H}^u_G g_{\hat{\mathbf{u}}}(t,x)) \\
&= \mathbf{A}^u V_{\hat{\mathbf{u}}}(t,x) - \mathbf{A}^u f_{\hat{\mathbf{u}}}(t,x,x) + \mathbf{A}^u f^x_{\hat{\mathbf{u}}}(t,x) - \mathbf{A}^u G \diamond g_{\hat{\mathbf{u}}}(t,x) + \mathbf{H}^u_G g_{\hat{\mathbf{u}}}(t,x).
\end{aligned}
$$

Recall that (t,x,u) was arbitrary. The above therefore implies that $\hat{\mathbf{u}}$ is maximal in (18), which is all we had left to prove. ∎

6 The general case

In this section we add a time-inconsistent *running payoff function* H and we also allow the time-inconsistent terminal payoff functions F and G to depend on the current time t. Specifically, we consider the following value function.

Definition 6.1 (Value function II)

$$
\begin{aligned}
J(t,x,\mathbf{u}) := E_{t,x}\left[\int_t^T H(r, X^{\mathbf{u}}_r, \mathbf{u}(r, X^{\mathbf{u}}_r), t, x) dr + F(t, X^{\mathbf{u}}_T, x)\right] \\
+ \qquad G(t, x, E_{t,x}\left[X^{\mathbf{u}}_T\right])
\end{aligned}
$$

where $F : [0,T] \times \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}$ *is continuous with* $|F(t,x,y)| \leq C_0(1+|x|^2)$ *for some constant* C_0 *which is independent of* t *(but which may depend on* y*),* $H : [0,T] \times \mathbb{R}^n \times U \times [0,T] \times \mathbb{R}^n \to \mathbb{R}$ *is continuous and bounded and* $G : [0,T] \times \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R}$ *satisfies* $G \in C^{1,2}([0,T) \times \mathbb{R}^n \times \mathbb{R}^n) \cap C([0,T] \times \mathbb{R}^n \times \mathbb{R}^n)$.

In this more general setting setting we consider the following definitions

$$
\begin{aligned}
&G \diamond g(t,x) := G(t,x,g(t,x)),\\
&\mathbf{H}^{\mathbf{u}}_G g(t,x) := G_y(t,x,g(t,x)) \cdot \mathbf{A}^{\mathbf{u}} g(t,x)\\
&f_{\mathbf{u}}(t,x,s,y) := E_{t,x}\left[\int_t^T H(r, X^{\mathbf{u}}_r, \mathbf{u}(r, X^{\mathbf{u}}_r), s, y)dr + F(s, X^{\mathbf{u}}_T, y)\right],
\end{aligned}
$$

where $G_y(x,y)$ is the gradient of G with respect y, for fixed (t,x). The definition $g_{\mathbf{u}}(t,x) := E_{t,x}\left[X^{\mathbf{u}}_T\right]$ is still the same.

It is easy to see that an analogous version of Lemma 2.4 holds in this more general case. Specifically, in this version of Lemma 2.4 we replace $f_{\mathbf{u}}(t,x,y)$ with $f_{\mathbf{u}}(t,x,s,y)$ and note that C_2 is independent also of s.

The equilibrium definition is analogous to Definition 3.2. The only difference is that the equilibrium quadruple is $(\hat{\mathbf{u}}, V_{\hat{\mathbf{u}}}(t,x), f_{\hat{\mathbf{u}}}(t,x,s,y), g_{\hat{\mathbf{u}}}(t,x))$, where we still let $V_{\hat{\mathbf{u}}}(t,x) := J(t,x,\hat{\mathbf{u}})$.

The extended HJB system is in this general case given in Definition 6.2, where $f(t,x,s,y) = f^{s,y}(t,x)$ and $\mathcal{A}^{\mathbf{u}}$ still operates only on variables in parenthesis.

Definition 6.2 (The extended HJB system II) *For $(t,x,s,y) \in [0,T) \times \mathbb{R}^n \times [0,T) \times \mathbb{R}^n$,*

$$
\begin{aligned}
\mathcal{A}^{\bar{\mathbf{u}}} f^{s,y}(t,x) + H(t,x,\bar{\mathbf{u}}(t,x),s,y) = 0, \;\; f^{s,y}(T,x) &= F(s,x,y)\\
\mathcal{A}^{\bar{\mathbf{u}}} g(t,x) = 0, \;\; g(T,x) &= x\\
\sup_{u\in U}\{\mathbf{A}^u V(t,x) - \mathbf{A}^u f(t,x,t,x) + \mathbf{A}^u f^{t,x}(t,x)&\\
-\mathbf{A}^u G \diamond g(t,x) + \mathbf{H}^u_G g(t,x) + H(t,x,u,t,x)\} &= 0\\
V(T,x) = F(T,x,x) + G(T,x,x)&
\end{aligned}
$$

where $\bar{\mathbf{u}}$ satisfies

$$
\begin{aligned}
\bar{\mathbf{u}}(t,x) \in \sup_{u\in U}\{&\mathbf{A}^u V(t,x) - \mathbf{A}^u f(t,x,t,x) + \mathbf{A}^u f^{t,x}(t,x)\\
&-\mathbf{A}^u G \diamond g(t,x) + \mathbf{H}^u_G g(t,x) + H(t,x,u,t,x)\} = 0.
\end{aligned}
$$

The definition of a regular equilibrium is almost the same as in Section 5. The difference is that Assumption 5.3 here regards the differentiability of $f^{s,y}_{\hat{\mathbf{u}}}(t,x)$ for fixed (s,y) and $f_{\hat{\mathbf{u}}}(t,x,t,x)$ instead of $f_{\hat{\mathbf{u}}}(t,x,x)$.

We are now ready to present the main result in the general case. The proof is very similar to the proof of Theorem 5.5 and is therefore omitted.

Theorem 6.3 *A regular equilibrium $(\hat{\mathbf{u}}, V(t,x), f_{\hat{\mathbf{u}}}(t,x,s,y), g_{\hat{\mathbf{u}}}(t,x))$ necessarily solves the extended HJB system (Definition 6.2). In particular, if a regular equilibrium exists then the extended HJB system has a solution.*

7 Example: a time-inconsistent quadratic regulator

A standard quadratic regulator problem

The linear-quadratic regulator is a classic problem in control theory. A typical stochastic linear-quadratic regulator problem in dimension $n = 1$ can be described as follows. The drift of the controlled process is linear and the volatility $\sigma > 0$ is a constant,

$$dX_s^{\mathbf{u}} = (c_1 X_s^{\mathbf{u}} + c_2 \mathbf{u}(s, X_s^{\mathbf{u}}))ds + \sigma dW_s, \quad X_t^{\mathbf{u}} = x.$$

The running and terminal payoffs are quadratic and correspond to the value function

$$E_{t,x}\left[\int_t^T (c_3 \mathbf{u}(s, X_s^{\mathbf{u}})^2 + c_4 (X_s^{\mathbf{u}})^2)ds + c_5 (X_T^{\mathbf{u}})^2\right]. \tag{24}$$

All c_i are constants and $c_3, c_4, c_5 > 0$. The optimization problem is to *minimize* the value function (24).

A time-inconsistent quadratic regulator

We consider a time-inconsistent version of the regulator problem which we call *a time-inconsistent quadratic regulator*. Let $n = 1$ and $U = [-a, a]$ for some constant $a > 0$. Let the terminal payoff F be given by $F(x, y) = (x - y)^2$ while the other payoffs G and H are identically equal to zero. Let $\sigma > 0$ be constant and $\mu(t, x, u) = -u^2$. It is easy to verify that this model satisfies the general conditions of Section 2. The resulting value function that the agents wish to *maximize* is

$$J(t, x, \mathbf{u}) = E_{t,x}\left[(X_T^{\mathbf{u}} - x)^2\right]. \tag{25}$$

The auxiliary functions are $g_{\mathbf{u}}(t, x) = E_{t,x}\left[X_T^{\mathbf{u}}\right]$ and

$$f_{\mathbf{u}}(t, x, y) = E_{t,x}\left[(X_T^{\mathbf{u}} - y)^2\right].$$

The dynamics are

$$dX_s^{\mathbf{u}} = -\mathbf{u}(s, X_s^{\mathbf{u}})^2 ds + \sigma dW_s, \quad X_t^{\mathbf{u}} = x. \tag{26}$$

Now, *guess* that $\hat{\mathbf{u}} = 0$ is an equilibrium control. Using (26), this implies that

$$\begin{aligned}
&f_{\hat{\mathbf{u}}}(t, x, y) = E_{t,x}\left[(X_T^{\hat{\mathbf{u}}} - y)^2\right] = E_{t,x}\left[(x + \sigma(W_T - W_t) - y)^2\right] \\
&= E_{t,x}\left[(x + \sigma(W_T - W_t))^2 - 2y(x + \sigma(W_T - W_t)) + y^2\right] \\
&= E_{t,x}\left[x^2 + 2x\sigma(W_T - W_t) + \sigma^2(W_T - W_t)^2\right] - 2xy + y^2 \\
&= x^2 + \sigma^2(T - t) - 2xy + y^2 = (x - y)^2 + \sigma^2(T - t).
\end{aligned}$$

Moreover, $g_{\hat{\mathbf{u}}}(t, x) = E_{t,x}\left[X_T^{\hat{\mathbf{u}}}\right] = E_{t,x}\left[x + \sigma(W_T - W_t)\right] = x$. This implies that,

$$\begin{gathered}
\frac{\partial f_{\hat{\mathbf{u}}}(t, x, y)}{\partial t} = -\sigma^2, \quad \frac{\partial f_{\hat{\mathbf{u}}}(t, x, y)}{\partial x} = 2x - 2y, \quad \frac{\partial^2 f_{\hat{\mathbf{u}}}(t, x, y)}{\partial x^2} = 2, \\
\frac{\partial g_{\hat{\mathbf{u}}}(t, x)}{\partial t} = 0, \quad \frac{\partial g_{\hat{\mathbf{u}}}(t, x)}{\partial x} = 1, \quad \frac{\partial^2 g_{\hat{\mathbf{u}}}(t, x)}{\partial x^2} = 0.
\end{gathered} \tag{27}$$

Remark 7.1 *We need the following general observation. Let (t, x) be an arbitrary point and let $\mathbf{u}$ be an arbitrary control. Consider a control $\mathbf{u}_{\theta_n}$ defined as being equal to $\mathbf{u}$ on the stochastic interval $[t, \theta_n]$ and equal to $\hat{\mathbf{u}}$ outside of this interval, where*

$$\theta_n := \inf\left\{s > t : (s, X_s^{\mathbf{u}}) \notin [t, t + h_n) \times B[x]\right\} \wedge T.$$

In the same way as in the proof of Theorem 5.5 it follows that $\mathbf{u}_{\theta_n}$ *is a control of the type defined in Definition 3.2, specifically* $\mathbf{u}_{\theta_n} = \mathbf{u}_{t+h_n}$*. Using the same arguments as in the proof of Theorem 5.5 we can, for any sufficiently differentiable function* $f_{\hat{\mathbf{u}}}(t,x,y)$*, show that*

$$f_{\hat{\mathbf{u}}}(t,x,x) - f_{\mathbf{u}_{\theta_n}}(t,x,x) = -E_{t,x}\left[\int_t^{\theta_n} \mathbf{A}^{\mathbf{u}(s,X_s^{\mathbf{u}})} f_{\hat{\mathbf{u}}}^x(s,X_s^{\mathbf{u}})ds\right]. \quad (28)$$

Let us return to our example and show that our guess is correct, by showing that the value function (25) satisfies the equilibrium condition (7) when we use the guessed control $\hat{\mathbf{u}} = 0$. Consider an arbitrary point (t,x) and an arbitrary control $\mathbf{u}$. Using that $n = 1$, σ is a constant, $\mu(t,x,u) = -u^2$ and (27) we see that

$$\begin{aligned}
&\mathbf{A}^{\mathbf{u}(s,X_s^{\mathbf{u}})} f_{\hat{\mathbf{u}}}^x(s,X_s^{\mathbf{u}}) \\
&= \frac{\partial f_{\hat{\mathbf{u}}}^x(s,X_s^{\mathbf{u}})}{\partial t} + \mu(s,X_s^{\mathbf{u}},\mathbf{u}(s,X_s^{\mathbf{u}}))\frac{\partial f_{\hat{\mathbf{u}}}^x(s,X_s^{\mathbf{u}})}{\partial x} + \frac{\sigma^2}{2}\frac{\partial^2 f_{\hat{\mathbf{u}}}^x(s,X_s^{\mathbf{u}})}{\partial x^2} \\
&= -\mathbf{u}(s,X_s^{\mathbf{u}})^2(2X_s^{\mathbf{u}} - 2x). \quad (29)
\end{aligned}$$

Now use the definition of $J(t,x,\mathbf{u})$, $G = 0$ with (6), (28) and (29), and $U = [-a,a]$, to obtain

$$\begin{aligned}
&J(t,x,\hat{\mathbf{u}}) - J(t,x,\mathbf{u}_{\theta_n}) \\
&= f_{\hat{\mathbf{u}}}(t,x,x) + G \diamond g_{\hat{\mathbf{u}}}(t,x) - (f_{\mathbf{u}_{\theta_n}}(t,x,x) + G \diamond g_{\mathbf{u}_{\theta_n}}) \\
&= E_{t,x}\left[\int_t^{\theta_n} \mathbf{u}(s,X_s^{\mathbf{u}})^2(2X_s^{\mathbf{u}} - 2x)ds\right] \\
&\geq -2a^2 E_{t,x}\left[\int_t^{\theta_n} |X_s^{\mathbf{u}} - x|ds\right]. \quad (30)
\end{aligned}$$

Arguments analogous to those in the proof of Theorem 5.5 imply that we may now use Lebesgue's differentiation theorem and dominated convergence as follows. Since $X^{\mathbf{u}}$ has continuous trajectories and $X_t^{\mathbf{u}} = x$, Lebesgue's differentiation theorem implies that $\frac{1}{h_n}\int_t^{\theta_n} |X_s^{\mathbf{u}} - x|ds$ converges to 0 as $n \to \infty$ for a.e. $\omega \in \Omega$. Using this and dominated convergence (note that $X_s^{\mathbf{u}}$ is bounded on $[t,\theta_n]$) we obtain from (30) that

$$\liminf_{n\to\infty} \frac{J(t,x,\hat{\mathbf{u}}) - J(t,x,\mathbf{u}_{\theta_n})}{h_n} \geq 0.$$

Since (t,x) and the alternative control $\mathbf{u}$ (and $B[x]$) were arbitrary it follows that $\hat{\mathbf{u}}$ is an equilibrium control (see Definition 3.2, and recall that $\mathbf{u}_{\theta_n} = \mathbf{u}_{t+h_n}$ and $\lim_{n\to\infty} h_n = 0$). Our guess was therefore correct and

$$(\hat{\mathbf{u}}, V_{\hat{\mathbf{u}}}(t,x), f_{\hat{\mathbf{u}}}(t,x,y), g_{\hat{\mathbf{u}}}(t,x)) = (0, \sigma^2(T-t), (x-y)^2 + \sigma^2(T-t), x) \quad (31)$$

is an equilibrium for the time-inconsistent quadratic regulator.

The corresponding extended HJB system

It is easy to see that our equilibrium (31) is regular (Definition 5.1): the equilibrium control $\hat{\mathbf{u}} = 0$ is right-continuous and the auxiliary and equilibrium value

functions $f_{\hat{\mathbf{u}}}(t,x,y) = (x-y)^2 + \sigma^2(T-t)$, $g_{\hat{\mathbf{u}}}(t,x) = x$ and $V_{\hat{\mathbf{u}}}(t,x) = \sigma^2(T-t)$ are sufficiently differentiable. Since the equilibrium is regular we may use Theorem 5.5 to draw the conclusion that it necessarily solves the corresponding extended HJB system. Let us verify, however, that this is the case.

The equilibrium (31) satisfies the boundary conditions of the extended HJB system (Definition 4.1), since $V_{\hat{\mathbf{u}}}(T,x) = \sigma^2(T-T) = 0 = (x-x)^2 = F(x,x) + G(x,x)$, $f_{\hat{\mathbf{u}}}(T,x,y) = (x-y)^2 + \sigma^2(T-T) = (x-y)^2 = F(x,y)$ and $g_{\hat{\mathbf{u}}}(T,x,y) = x$, where used that $G(x,y) = 0$.

Note that $G(x,y) = 0$ also implies that $\mathbf{A}^u G \diamond g_{\hat{\mathbf{u}}}(t,x) = \mathbf{H}^u_G g_{\hat{\mathbf{u}}}(t,x) = 0$. Moreover, $V_{\hat{\mathbf{u}}}(t,x) = \sigma^2(T-t) = f_{\hat{\mathbf{u}}}(t,x,x)$, which implies that $\mathbf{A}^u V_{\hat{\mathbf{u}}}(t,x) = \mathbf{A}^u f_{\hat{\mathbf{u}}}(t,x,x)$. With μ and σ defined as above, it therefore follows that we now only need to show that the following equations hold in order to prove that the equilibrium (31) solves the extended HJB system

$$\frac{\partial f^y_{\hat{\mathbf{u}}}(t,x)}{\partial t} - \hat{\mathbf{u}}(t,x)^2 \frac{\partial f^y_{\hat{\mathbf{u}}}(t,x)}{\partial x} + \frac{1}{2}\sigma^2 \frac{\partial^2 f^y_{\hat{\mathbf{u}}}(t,x)}{\partial x^2} = 0 \tag{32}$$

$$\frac{\partial g_{\hat{\mathbf{u}}}(t,x)}{\partial t} - \hat{\mathbf{u}}(t,x)^2 \frac{\partial g_{\hat{\mathbf{u}}}(t,x)}{\partial x} + \frac{1}{2}\sigma^2 \frac{\partial^2 g_{\hat{\mathbf{u}}}(t,x)}{\partial x^2} = 0 \tag{33}$$

$$\sup_{u\in U}\left\{ \frac{\partial f^x_{\hat{\mathbf{u}}}(t,x)}{\partial t} - u^2 \frac{\partial f^x_{\hat{\mathbf{u}}}(t,x)}{\partial x} + \frac{1}{2}\sigma^2 \frac{\partial^2 f^x_{\hat{\mathbf{u}}}(t,x)}{\partial x^2} \right\} = 0 \tag{34}$$

$$\hat{\mathbf{u}}(t,x) \in \arg\sup_{u\in U}\left\{ \frac{\partial f^x_{\hat{\mathbf{u}}}(t,x)}{\partial t} - u^2 \frac{\partial f^x_{\hat{\mathbf{u}}}(t,x)}{\partial x} + \frac{1}{2}\sigma^2 \frac{\partial^2 f^x_{\hat{\mathbf{u}}}(t,x)}{\partial x^2} \right\} \tag{35}$$

where (32) and (33) are what remains of the Kolmogorov part, and (34) and (35) are what remains of the HJB part. The derivatives of the auxiliary function $f_{\hat{\mathbf{u}}}(t,x,y) = (x-y)^2 + \sigma^2(T-t)$ are given in (27). Using these we write the left side of (32) as

$$-\sigma^2 - \hat{\mathbf{u}}(t,x)^2(2x-2y) + \frac{1}{2}\sigma^2 2 = -\hat{\mathbf{u}}(t,x)^2(2x-2y)$$

which clearly is equal to zero, since the equilibrium control is $\hat{\mathbf{u}} = 0$. This verifies that the equilibrium (31) solves (32). We can verify that the equilibrium (31) solves (33) in the same way. Similarly, (34) can be written as

$$\sup_{u\in U}\left\{ -\sigma^2 - u^2(2x-2x) + \frac{1}{2}\sigma^2 2 \right\} = 0$$

where $u = 0$ clearly attains the supremum 0 in the left side. This means that (34) and (35) are satisfied by $f_{\hat{\mathbf{u}}}(t,x,y) = (x-y)^2 + \sigma(T-t)$ and $\hat{\mathbf{u}} = 0$.

We have thus verified that the equilibrium (31) solves the corresponding extended HJB system.

A time-consistent version of the time-inconsistent regulator

The time-inconsistent quadratic regulator problem studied above is genuinely time-inconsistent in the sense that the equilibrium control $\hat{\mathbf{u}} = 0$ is not an optimal control in following time-consistent version of this problem: Consider the time-inconsistent quadratic regulator problem, but now let the agent at the initial point $(0, x_0)$ be able to dictate which control $\mathbf{u}$ should be used until T. This is clearly a standard stochastic control problem with optimal value function

$$\sup_{\mathbf{u}\in\mathbf{U}} E_{t,x}[(X^{\mathbf{u}}_T - x_0)^2]. \tag{36}$$

Therefore, if $\mathbf{u} = 0$ were an optimal control then the optimal value function, denote it by K, would be given by (cf. the calculations above (27))

$$K(t,x) = E_{t,x}[(X_T^{\mathbf{u}} - x_0)^2] = (x - x_0)^2 + \sigma^2(T-t). \tag{37}$$

Therefore, if $\mathbf{u} = 0$ were an optimal control then we would be able to use the standard (time-consistent) stochastic control version of Theorem 5.5 (see e.g. [20, ch. 11]) to deduce that the function K in (37) would solve the standard HJB corresponding to (36) (and (26)),

$$\sup_{u\in U}\left\{\frac{\partial K(t,x)}{\partial t} - u^2\frac{\partial K(t,x)}{\partial x} + \frac{1}{2}\sigma^2\frac{\partial^2 K(t,x)}{\partial x^2}\right\} = 0, \quad K(T,x) = (x-x_0)^2. \tag{38}$$

It is easy to verify that the function K in (37) *does not* solve (38) which therefore implies that $\mathbf{u} = 0$ cannot be an optimal control for this time-consistent version of our problem.

Remark 7.2 *Another time-inconsistent version of the regulator problem is studied in [2].*

8 Conclusions and remarks

We study time-inconsistent stochastic control. Our model is that of a general (Markovian) Itô diffusion and general time-inconsistent terminal payoff functions, see Section 2. We also study a more general case including a time-inconsistent running payoff function, see Section 6. The standard notion of optimality is not suitable for time-inconsistent optimization and, in Section 3, we therefore define the time-inconsistent stochastic control *equilibrium* (found also in [2, 3, 4, 5, 11, 12]). In Section 4, we define a system of PDEs called *the extended HJB system* (proposed in [2]). Our main result is Theorem 5.5 (and the more general Theorem 6.3), which says that a regular equilibrium (*if it exists*) is necessarily a solution to the corresponding extended HJB system. The proof is based on probabilistic methods.

The remarks below reveal a potential for future research in time-inconsistent stochastic control.

Remark 8.1 (Existence of equilibrium controls,) *The problem of finding general model conditions (i.e. for σ, μ, U, F, G and H) which guarantee the existence of an equilibrium control, and a solution to the extended HJB system, is still open. For related remarks see [2, section 4.2] and [5, section 6].*

Remark 8.2 (Differentiability) *For any particular model one should not a priori expect the differentiability in Assumption 5.3 to be satisfied. It involves a lot of work to find useful model conditions such that similar assumptions are satisfied for standard (time-consistent) stochastic control (for such conditions see e.g. [13, 14, 19, 24]). Indeed, Fleming and Rishel (2012) [13, p. 154] write the following about a similar differentiability assumption in standard stochastic control:* "Substantial difficulties are encountered if one seeks to put all of the above formal discussion on a precise basis. It has been implicitly assumed that $W^0(t,\cdot)$ *[the optimal value function]* is in the domain of $\mathbf{A}^{\mathbf{u}}(t)$ *[the differential operator]*, which is not always the case."

We also remark that we would clearly have to impose more regularity for the variable y in the terminal payoff function F in (4) in order for Assumption 5.3 to be reasonable.

Remark 8.3 (On the possibility of several equilibria) *In game theory there is generally no reason to expect anything else than that there may exist several equilibria, or no equilibrium. Note, however, that the investigations in the present paper regard any fixed equilibrium control $\hat{\mathbf{u}}$, and if, in a particular case, there should be more than one equilibrium control, then our results are true for any fixed equilibrium control. Moreover, we do not expect that different equilibrium controls necessarily correspond to different equilibrium value functions, although they may do so in particular cases.*

Remark 8.4 (Viscosity solutions) *For the standard time-consistent stochastic control theory, it is well-known that rather strong conditions for the model are needed in order for the value function to be sufficiently differentiable to be able to solve the HJB in the classical sense. This is true also when considering a.e. differentiability. Clearly, there is no reason to expect anything else for time-inconsistent stochastic control. For the standard time-consistent stochastic control theory, this problem was famously solved by the introduction of a type of weak solutions to PDEs known as* viscosity solutions*, see [7]. Viscosity solutions for extended HJB systems have so far not been studied.*

Remark 8.5 (More examples) *Several particular time-inconsistent stochastic control problems studied in a mathematical way can be found in e.g. [2, 3, 5, 6, 8, 11, 21].*

Acknowledgments

The author is grateful to Tomas Björk and Jan-Erik Björk for helpful discussions.

References

[1] R. F. Bass. The measurability of hitting times. *Electron. Comm. Prob*, 15:99–105, 2010.

[2] T. Björk, M. Khapko, and A. Murgoci. On time inconsistent stochastic control in continuous time. *Forthcoming in Finance and Stochastics, (SSRN preprint 2890162)*, 2016.

[3] T. Björk, M. Khapko, and A. Murgoci. Time inconsistent stochastic control in continuous time: Theory and examples. arXiv preprint 1612.03650, 2016.

[4] T. Björk and A. Murgoci. A general theory of Markovian time inconsistent stochastic control problems. SSRN preprint 1694759, 2010.

[5] T. Björk and A. Murgoci. A theory of Markovian time-inconsistent stochastic control in discrete time. *Finance and Stochastics*, 18(3):545–592, 2014.

[6] T. Björk, A. Murgoci, and X. Y. Zhou. Mean-variance portfolio optimization with state-dependent risk aversion. *Mathematical Finance*, 24(1):1467–9965, 2014.

[7] M. G. Crandall, H. Ishii, and P.-L. Lions. User's guide to viscosity solutions of second order partial differential equations. *Bulletin of the American Mathematical Society*, 27(1):1–67, 1992.

[8] C. Czichowsky. Time-consistent mean-variance portfolio selection in discrete and continuous time. *Finance and Stochastics*, 17(2):227–271, 2013.

[9] D.-M. Dang, P. A. Forsyth, and Y. Li. Convergence of the embedded mean-variance optimal points with discrete sampling. *Numerische Mathematik*, 132(2):271–302, 2016.

[10] B. Djehiche and M. Huang. A characterization of sub-game perfect equilibria for SDEs of mean-field type. *Dynamic Games and Applications*, 6(1):1–27, 2015.

[11] I. Ekeland and A. Lazrak. Being serious about non-commitment: subgame perfect equilibrium in continuous time. arXiv preprint math/0604264, 2006.

[12] I. Ekeland and T. Pirvu. Investment and consumption without commitment. *Mathematics and Financial Economics*, 2(1):57–86, 2008.

[13] W. H. Fleming and R. W. Rishel. *Deterministic and stochastic optimal control*, volume 1. Springer, 2012.

[14] W. H. Fleming and H. M. Soner. *Controlled Markov processes and viscosity solutions, 2nd edition*. Springer, 2006.

[15] S. M. Goldman. Consistent plans. *The Review of Economic Studies*, 47(3):533–537, 1980.

[16] I. Gyöngy and N. Krylov. Existence of strong solutions for Itô's stochastic equations via approximations. *Probability theory and related fields*, 105(2):143–158, 1996.

[17] I. Karatzas and S. E. Shreve. *Brownian Motion and Stochastic Calculus (Graduate Texts in Mathematics), 2nd edition*. Springer, 1991.

[18] K. Krusell and A. Smith. Consumption-savings decisions with quasi-geometric discounting. *Econometrica*, 71(1):365–375, 2003.

[19] N. V. Krylov. *Controlled diffusion processes*, volume 14. Springer, 2008.

[20] B. Øksendal. *Stochastic Differential Equations: An Introduction with Applications (Universitext)*. Springer, 6 edition, 2003.

[21] J. L. Pedersen and G. Peskir. Optimal mean-variance selling strategies. *Mathematics and Financial Economics*, 10(2):203–220, 2016.

[22] B. Peleg and M. E. Yaari. On the existence of a consistent course of action when tastes are changing. *The Review of Economic Studies*, 40(3):391–401, 1973.

[23] R. Strotz. Myopia and inconsistency in dynamic utility maximization. *The Review of Economic Studies*, 23(3):165–180, 1955.

[24] N. Touzi. Stochastic control problems, viscosity solutions and application to finance. Technical report, Special research semester on Financial Markets, Scuola Normale Superiore, Pisa, 2004.

[25] A. J. Veretennikov. On strong solutions and explicit formulas for solutions of stochastic integral equations. *Math. USSR Sbornik*, 39(3):387–403, 1981.

[26] N. Vieille and J. W. Weibull. Multiple solutions under quasi-exponential discounting. *Economic Theory*, 39(3):513–526, 2009.

[27] H. Wang and Z. Wu. Time-inconsistent optimal control problem with random coefficients and stochastic equilibrium HJB equation. *Mathematical control and related fields*, 5(3):651–678, 2015.

[28] J. Yong. Deterministic time-inconsistent optimal control problems - an essentially cooperative approach. *Acta Mathematicae Applicatae Sinica*, 28(1):1–30, 2012.

[29] J. Yong. Time-inconsistent optimal control problems and the equilibrium HJB equation. *Mathematical Control and Related Fields*, 2(3):271–329, 2012.